





55 year-old female with history of bladder cancer; presents with dysuria during a visit to her primary care physician. Given her history and symptoms, physician calls for a BTA stat test and patient is referred to a urologist.



52 year-old male, routinely monitored for recurrent bladder cancer. Physician orders a BTA stat test which later returns a positive result. Cytology is negative. Cystoscopy detects a tumor, confirming the BTA stat test result.

73 year-old male with a low-grade TCC detected and treated two years ago; found positive for microhematuria in a routine laboratory urinalysis. Given the low-grade nature of the initial tumor, physician uses the BTA stat test instead of cytology.



60 year-old male with TCC recurrence detected and treated three years ago. History of recurrence calls for a vigilant detection program. Surveillance is enhanced by using BTA stat tests between scheduled cystoscopies.

Detection of Recurrent Bladder Cancer in Only 5 Minutes

... with the BTA stat® Test

Bladder Cancer: A Common Disease That Requires Lifelong Management

Bladder cancer is the fourth most common cancer among men

One in five patients diagnosed with bladder cancer dies from the disease

Bladder cancer recurs in over 50% of patients calling for regular testing and lifelong management

Early Detection is the Key to Bladder Cancer Survival

Risk Factors For Bladder Cancer

Tobacco smokers over 50 years of age

Persons who have worked in the chemical, dye, metal, rubber, or leather tanning industries


Previous diagnosis of bladder cancer

Bladder Cancer Surveillance

The true clinical partner to Cystoscopy

Simple collection allows for improved patient comfort and compliance

Bladder Cancer 5 Year Survival Rate



Increased detection rates may result in fewer cancers missed and improved outcomes.

The BTA stat[®] Test

... aid to the management of bladder cancer patients

An Advanced Method Delivers Objective Results in Only Five Minutes


Uses monoclonal antibodies to detect the presence of bladder tumor associated antigen in urine

Detects early stage and grade cancers that cytology often misses

Returns fast, objective result - a line in the patient window indicates a positive result

Requires only one voided urine sample, with no sample preparation

Adjunct test to cystoscopy



Ordering Information

661210 - 10 Test Kit

661230 - 30 Test Kit

661203 - Control Kit, 15 Determinations

Distributed by: **CLIAwaived.com[™]**

San Diego, CA 92121

tel 858-481-5031


toll free 888-882-7739

www.cliawaived.com

The BTA stat test offers accurate, objective test results with outstanding sensitivity and specificity


The BTA stat Test vs Cytology

Sensitivity by Grade



The BTA stat Test Specificity

For Healthy Subjects
Age Range 35-86



Non-genitourinary
diseases and cancers

